

Scaling approach to the phase diagram of quantum Hall systems

M. O. GOERBIG^{1,2} AND C. MORAIS SMITH¹

¹ *Département de Physique, Université de Fribourg, Pérolles, CH-1700 Fribourg, Switzerland.*

² *Laboratoire de Physique des Solides, Université Paris-Sud, F-91405 Orsay, France.*

PACS. 73.43.-f – Quantum Hall effects.

PACS. 73.43.Nq – Quantum phase transitions.

PACS. 73.20.Qt – Electron Solids.

Abstract. – We present a simple classification of the different liquid and solid phases of quantum Hall systems in the regime where the Coulomb interaction between electrons is significant, i.e. away from integral filling factors. This classification, and a criterion for the validity of the mean-field approximation in the charge-density-wave phase, is based on scaling arguments concerning the effective interaction potential of electrons restricted to an arbitrary Landau level. Finite-temperature effects are investigated within the same formalism, and a good agreement with recent experiments is obtained.

Two-dimensional electron systems (2DES) in a perpendicular magnetic field exhibit a rich variety of phases, ranging from incompressible quantum liquids, which are responsible for the integral and fractional quantum Hall effects (IQHE and FQHE), to electron-solid phases such as charge density waves (CDWs) and the Wigner crystal (WC). The IQHE is found when the electron density n_{el} is an integral multiple of the density of states per Landau level (LL) $n_B = 1/2\pi l_B^2$, where $l_B = \sqrt{\hbar/eB}$ is the magnetic length [1], and can be described within a single-particle picture. On the other hand, when the filling factor $\nu = n_{el}/n_B$ is non-integral, the highest filled LL has only a partial filling $\bar{\nu} = \nu - n$, and it becomes essential to include the Coulomb interaction [2]. It lifts the LL degeneracy and leads to a rich phase diagram. At extremely low electron densities, an insulating phase has been observed whose properties are attributed to WC formation [3]. At $\nu = p/q$, with p, q integral and q odd, the FQHE is observed in the two lowest LLs [4]. The corresponding ground state is an incompressible liquid, analogous to the IQHE if described in terms of composite fermions (CFs) [5]. At $\bar{\nu} = 1/2$ in higher LLs, huge anisotropies have been detected in the longitudinal magnetoresistance, indicating the formation of a unidirectional CDW [6]. From the theoretical point of view, Hartree-Fock calculations predict correctly a CDW formation around $\bar{\nu} = 1/2$ in higher LLs [7, 8], but have failed to describe the FQHE regime. In this letter we propose a new scaling approach, which classifies the liquid and solid phases observed in quantum Hall systems in terms of length scales, and remains valid in all LLs. In addition, we clarify the reasons for breakdown of the Hartree-Fock approximation in the lowest LLs and establish a

criterion for the appearance of recently observed FQHE features in the WC regime at finite temperatures [9].

Because in the high-magnetic-field limit the Coulomb interaction between the electrons constitutes a smaller energy scale than the gap between LLs, inter-LL excitations are high-energy degrees of freedom. At integral filling, they are the only excitations, but may be neglected when considering a system at $\nu \neq n$ because in this case low-energy excitations become possible within the same LL [10]. The kinetic energy of the electrons may therefore be set to zero, and the Coulomb interaction remains as the only energy scale in the problem if the electrostatic potential due to underlying impurities is small. One obtains a system of strongly correlated electrons described by the Hamiltonian

$$\hat{H}_n = \frac{1}{2} \sum_{\mathbf{q}} v(q) [F_n(q)]^2 \bar{\rho}(-\mathbf{q}) \bar{\rho}(\mathbf{q}), \quad (1)$$

where $v(q) = 2\pi e^2/\epsilon q$ is the two-dimensional Fourier transform of the Coulomb interaction. In this model, one considers only interactions between spinless electrons within the n th LL described by the density operators $\langle \rho(\mathbf{q}) \rangle_n = F_n(q) \bar{\rho}(\mathbf{q})$, where $\rho(\mathbf{q})$ is the usual electron density in reciprocal space. The factors $F_n(q) = L_n(q^2/2) \exp(-q^2/4)$, with the Laguerre polynomials $L_n(x)$, arise from the wave functions of electrons in the n th LL and may be absorbed into an effective interaction potential $v_n(q) = v(q)[F_n(q)]^2$. The quantum nature of the problem is contained in unusual commutation relations for the electron density operators [11],

$$[\bar{\rho}(\mathbf{q}), \bar{\rho}(\mathbf{k})] = 2i \sin \left(\frac{(\mathbf{q} \times \mathbf{k})_z l_B^2}{2} \right) \bar{\rho}(\mathbf{q} + \mathbf{k}). \quad (2)$$

The Hamiltonian (1) together with the commutation relations (2) defines the full model, which was used recently as a starting point for the description of the FQHE in the lowest LL [12] and for the formation of CDWs in higher LLs [7, 8]. Note the formal equivalence between electrons in the lowest LL at $\nu = \nu_0$ and electrons in a higher LL at $\bar{\nu} = \nu_0$ within this model.

Deeper insight into the structure of the model is obtained by transforming the effective interaction potential back to real space. In appropriate units, one may derive a universal scaling function $\tilde{v}(r)$,

$$v_n(r) = \sum_{\mathbf{q}} e^{-i\mathbf{q} \cdot \mathbf{r}} v_n(q) \approx \frac{\tilde{v}(r/R_C)}{\sqrt{2n+1}}, \quad (3)$$

where $R_C = l_B \sqrt{2n+1}$ is the cyclotron radius. This scaling form becomes exact in the limit $n \rightarrow \infty$ because then $F_n(q) \rightarrow J_0(q\sqrt{2n+1})$. However, it is valid also at low values of n as can be seen from fig. 1, where the rescaled results of $v_n(r)$ are shown for the LLs $n = 1, \dots, 5$. The universal function exhibits a plateau of width $2R_C$ superimposed on the bare $1/r$ Coulomb potential, which is retrieved at large distances. Although the bare Coulomb potential possesses no characteristic length scale, a component whose range is characterised by R_C is introduced in the effective interaction potential. This permits a classification of the different phases according to the ratio of the average electronic separation $d \sim l_B/\sqrt{\bar{\nu}}$ in the n th LL and the range $2R_C$ of the effective interaction.

Quasi-classical limit (WC): $d \gg 2R_C$. In the limit of low density, the average distance between electrons interacting via the $1/r$ Coulomb potential is much larger than the spatial extent R_C of their wave functions. Quantum corrections to the classical result are of order $\mathcal{O}(R_C/d)$ and may thus be neglected. Classically, the electrons are arranged in a triangular WC in order to minimise the repulsive Coulomb interaction. The transition line which separates the WC phase from other phases is obtained by comparing d and R_C as functions




Fig. 1 – Effective interaction potential in real space. Points correspond to $n = 1, 2, \dots, 5$. Gray line: pure Coulomb potential.

of $\bar{\nu}$ and n (gray line in fig. 2). It is given by $\bar{\nu}_n^{WC} = \nu_0^{WC}/(2n+1)$, where ν_0^{WC} is the critical filling factor below which the WC is found in the lowest LL. Theoretical calculations predict $\nu_0^{WC} \approx 1/6.5$ for clean samples [13], while experimentally the onset of WC behaviour is observed around $\nu = 1/5$ [3].

Electrical transport in the WC phase arises either by a collective sliding mode or by the propagation of crystal dislocations. Sliding is suppressed by pinning of the WC due to residual impurities in the sample, and the number of dislocations is reduced by lowering the temperature. The experimental evidence for a WC phase arises from transport measurements [3], which indicate an insulating phase. In principle, this insulating behaviour could be attributed also to the localization of electrons by impurities [14]. However, because the samples used for the measurements are extremely clean, this is unlikely to be the case.

Mean-field limit (CDWs): $d \ll 2R_C$. Within the effective-potential framework, an arbitrarily chosen particle interacts strongly with a number of neighbours which can be estimated as $N_{n.n.} \approx \pi(2R_C)^2 \bar{n}_{el} = 2(2n+1)\bar{\nu}$, where $\bar{n}_{el}$ is the density of electrons in the n th LL. A mean-field approximation, such as Hartree-Fock, is valid for large $N_{n.n.}$, where each particle may be considered to interact only with an averaged background, without being influenced by the individual motions of its neighbours. This limit cannot be obtained in the lowest LL, where the cyclotron radius coincides with the magnetic length l_B . The average electronic separation would have to be shorter than this length, which constitutes the smallest possible spacing because each electronic state occupies a minimal surface $\sigma = 2\pi l_B^2$. We note in addition that in the lowest LL the effective interaction potential does not exhibit a plateau as for $n \geq 1$. The dark gray line in fig. 2 shows the relation $\bar{\nu}_n^{CDW} = N_c/(2n+1)$. Comparison with experiment yields $N_{n.n.} = N_c \sim 5$ for the limiting value above which the mean-field approximation is justified [6].

The validity of the mean-field approximation in the regime $d \ll 2R_C$ also becomes apparent in Fourier space, where the wave vectors are renormalised in the same manner as the distances in eq. (3). At large n , only the small-wave-vector limit remains important, and the commutation relation (2) for the density operators becomes

$$\left[\bar{\rho} \left(\frac{\mathbf{q}}{\sqrt{2n+1}} \right), \bar{\rho} \left(\frac{\mathbf{k}}{\sqrt{2n+1}} \right) \right] \approx i \frac{(\mathbf{q} \times \mathbf{k})_z l_B^2}{2n+1} \bar{\rho} \left(\frac{\mathbf{q} + \mathbf{k}}{\sqrt{2n+1}} \right) = \mathcal{O} \left(\frac{1}{n} \right),$$




Fig. 2 – Phase diagram at $T = 0$. Vertical and horizontal axes indicate, respectively, the LL index n and the partial filling $\bar{\nu}$. For completely filled levels $\bar{\nu} = 1$ one observes the IQHE (gray circles). The phase diagram is only shown up to half-filling $\bar{\nu} = 1/2$ because of particle-hole symmetry. Black circles denote points at which the FQHE is found in experiments, the thick gray line shows the region of an experimentally observed insulating phase attributed to WC formation [3], and gray squares indicate points where the stripe phase is observed [6]. The full lines represent the transition between different phases and move towards the FQHE liquid phase in the presence of impurities (see arrows and broken lines).

after rescaling and expansion of the sine function. The complicated algebraic structure of the density operators, and thus the quantum mechanical nature of the problem, become less important in higher LLs (larger n). This quasi-classical limit is different from the WC regime, in which the overlap of different electronic wave functions may be neglected. Here their overlap is sufficiently strong that the exchange interaction, which is included at the mean-field level, is essential.

The mean-field solution of the Hamiltonian (1) reveals that the ground state in this limit is a CDW with a characteristic period on the order of the cyclotron radius R_C [7, 8]. This clustering of electrons, in spite of their repulsive interaction, may be understood qualitatively from the form of the effective interaction potential in fig. 1: if two electrons approach more than their average separation d , only a small additional energy cost is incurred because of the plateau in the region $r < 2R_C$. However, a large energy on the order of the height of the plateau may be gained if the clustered electrons thus reduce the number of other electrons with which they interact strongly. The boundary above which the mean-field approximation becomes valid need not necessarily coincide with the CDW-FQHE phase transition. A detailed calculation of the ground-state energy would be needed to determine the exact transition line [7, 8, 15], but the present scaling investigations serve as an upper limit for this boundary.

In the region $\bar{\nu} < 1/2$, charged clusters (or “bubbles”) of several electrons form a super-WC to minimise their Coulomb repulsion (triangular CDW). Exactly at $\bar{\nu} = 1/2$, however, these bubbles percolate to form lines and thus give rise to a “stripe” pattern (unidirectional CDW). This breaking of rotational symmetry arises because of a competing symmetry: the particle-hole symmetry becomes exact at half filling. A phase of bubbles would violate this symmetry, and a very small residual anisotropy in the underlying crystal suffices to fix the direction of the stripes [16]. However, there are other inhomogeneous charge configurations, which would

also account for these symmetries, and there is theoretical evidence that a stripe pattern becomes unstable to the formation of an anisotropic WC at very low temperatures, breaking the particle-hole symmetry [17]. The experimental evidence for stripe phases in quantum Hall systems consists of very large anisotropies in the magnetoresistance with respect to two orthogonal directions [6]. This suggests an interpretation in terms of charge stripes, where easy electron transport along the stripe edges leads to a small resistance, but transport across the stripes involves relatively rare tunneling processes, thus explaining the large resistance in the orthogonal direction [17,18].

Quantum limit (FQHE): $d \sim 2R_C$. In the regime of the phase diagram intermediate between the WC and CDW phases, the FQHE is observed in the lowest two LLs [4]. Recent theoretical and experimental studies [12,19] support the description of the FQHE in terms of an incompressible liquid of CFs, each of which consists of a bound state of an electron and a vortex-like collective excitation which carries a charge of opposite sign [5]. The scaling arguments presented above show that in the quantum limit only one length scale is present in the problem, and thus no perturbative approaches are applicable.

Effect of impurities and finite temperature. The arguments presented above suggest that the observation of the FQHE is possible in higher LLs, e.g. at $\bar{\nu} = 1/5$ for $n = 2$. However, this phase has not yet been observed, and some experiments even suggest a CDW ground state in this regime [20]. Failure to observe the FQHE in higher LLs is likely to be due to the residual impurities in the samples, which favour crystalline structures such as the WC and CDWs [3]. Deformation of the inhomogeneous charge structure makes these phases better adapted to follow an underlying electrostatic potential than is an incompressible, homogeneous liquid. The two transition lines therefore move towards the quantum-liquid phase, as shown by the arrows and broken lines in fig. 2. The region where the FQHE is observed can become extremely narrow in higher LLs, and may even vanish, leading to a direct crossover between the triangular CDW and WC phases with no quantum melting in the intermediate regime $d \sim 2R_C$. However, samples of higher mobility and even lower impurity concentrations are expected to reveal FQHE states in the LL $n = 2$ in the future [21].

So far we have discussed the different phases determined by the two length scales d and R_C only at $T = 0$. Finite temperatures will introduce an additional thermal length scale defined by comparing the thermal energy $k_B T$ and the Coulomb interaction between electrons. One thus obtains $l_T = e^2 / \epsilon k_B T$, which corresponds to the de Broglie wavelength of the free electron gas.

If $l_T \gg R_C$, i.e. at low temperatures, thermal fluctuations may be neglected compared to quantum fluctuations of the correlated electron liquid. At $l_T \sim R_C$ thermal fluctuations destroy the quantum correlations and the FQHE disappears. In the mean-field limit, local crystalline structure of the CDW vanishes when thermal fluctuations become important on the length scale of the CDW periodicity $l_T \sim R_C$. This leads to an estimate of the CDW melting temperature $T_{CDW}(n) = C e^2 / \epsilon k_B l_B \sqrt{2n+1}$, with C a dimensionless constant, in agreement with previous work [7]. Because CDW states in high LLs are observed at relatively low magnetic fields (decreasing LL separation), inter-LL excitations have to be included, giving rise to a screening of the Coulomb interaction. The dielectric constant ϵ may then be replaced by $\epsilon(n) \sim 2n+1$ [7,10]. We stress that these scaling arguments provide an estimate of the melting temperature for *local* CDW order. The anisotropy observed at half-filling in higher LLs [6] vanishes at lower temperatures, indicating an isotropic distribution of local stripes, as proposed in a liquid-crystal picture [22].

Finite-temperature effects are most complex in the WC phase. Minima in the longitudinal magnetoresistance, similar to the ones arising in the FQHE regime, are experimentally observed above a temperature T_1 at lowest-LL filling fractions, where the WC phase is ex-

pected at $T = 0$ [9]. These minima vanish above a second temperature $T_2 > T_1$. An estimate for T_1 can be obtained from the Lindemann criterion [23]: the WC melts when the average displacement of an electron due to thermal fluctuations is a substantial fraction of the lattice constant $\langle \Delta r^2 \rangle = c_L^2 d^2$ with $c_L \sim 0.1$. Equating the potential energy for a small displacement of an electron in a WC, $e^2 \langle \Delta r^2 \rangle / \epsilon d^3$, to the thermal energy yields a melting temperature $T_1 \approx c_L^2 e^2 / \epsilon k_B d$, which is independent of the magnetic field, in good agreement with recent experiments [9]. This temperature corresponds to a thermal length $l_T \sim d / c_L^2 \gg l_B$. The liquid phase therefore exhibits quantum coherence on a length scale l_T , and may be described locally by the Laughlin wave function for temperatures such that $l_T > l_B$, thus displaying features of the FQHE. The scaling estimate $l_T \sim l_B$ for the definition of a temperature T_2 at which this coherence is lost turns out, however, to be rather crude because it neglects a renormalization of the magnetic length due to CF formation in the quantum limit. In the CF picture, the minima disappear when the temperature reaches the activation gap [12]. This leads to the relation $T_2 \approx T_C / (2ps \pm 1)$, where T_C is a constant, and the integers p, s are related to the filling factor $\bar{\nu} = p / (2ps \pm 1)$ for the FQHE states. Comparison with experiment [9] suggests a value $T_C \sim 2\text{K}$ around $\nu = 1/6$.

Conclusions. We have discussed the different solid and liquid phases of quantum Hall systems using straightforward scaling arguments. Although the bare Coulomb potential is scale-free, the effective potential of electrons restricted to the n th LL has a range of strong interaction characterised by the cyclotron radius. The ratio between this range and the average separation d of the electrons classifies the different phases at $T = 0$. At finite temperature, a further length scale l_T enters. For $d \gg 2R_C$ a WC is formed, which melts at a temperature T_1 into a quantum liquid showing features of FQHE states. Our estimates for the melting temperature T_1 show that it does not vary with B , in agreement with recent experiments [9]. The quantum coherence of the electrons is not destroyed until a higher temperature T_2 . In the opposite limit $d \ll 2R_C$ the mean-field approximation is justified and predicts a CDW ground state. We present a criterion for the validity of this approximation, which excludes CDW formation in the lowest LLs, where the required condition cannot be satisfied because R_C coincides with the smallest length l_B in the system. In an intermediate regime $d \sim 2R_C$ quantum melting of the crystalline structures leads to a liquid phase, which is incompressible at certain filling factors where the FQHE is observed. The presence of impurities reduces the FQHE regime in the phase diagram because solid phases are better adapted to follow an underlying impurity potential.

* * *

We acknowledge fruitful discussions with D. Baeriswyl, L. Benfatto, K. Borejsza, P. Lederer, R. Morf, B. Normand, V. Pasquier, and J. Wakeling. This work was supported by the Swiss National Foundation for Scientific Research under grant No. 620-62868.00.

REFERENCES

- [1] V. KLITZING K. *et al.*, *Phys. Rev. Lett.*, **45** (1980) 494.
- [2] If one takes into account the Zeeman splitting of each LL, the partial filling factors of the lower and upper spin branches of the n th LL are, respectively, $\bar{\nu} = \nu - 2n$ and $\bar{\nu} = \nu - (2n + 1)$.
- [3] JIANG H. *et al.*, *Phys. Rev. Lett.*, **65** (1990) 633; *Phys. Rev. B*, **44** (1991) 8107; ENGEL L. W. *et al.*, *Solid State Commun.*, **104** (1997) 167.
- [4] TSUI D. C. *et al.*, *Phys. Rev. Lett.*, **48** (1982) 1559; LAUGHLIN R. B., *Phys. Rev. Lett.*, **50** (1983) 1395.
- [5] JAIN J. K., *Phys. Rev. Lett.*, **63** (1989) 199; *Phys. Rev. B*, **41** (1990) 7653; *Composite Fermions*, edited by HEINONEN O. (World Scientific) 1998.

- [6] LILLY M. P. *et al.*, *Phys. Rev. Lett.*, **82** (1999) 394; DU R. R. *et al.*, *Solid State Commun.*, **109** (1999) 389.
- [7] KOULAKOV A. A. *et al.*, *Phys. Rev. Lett.*, **76** (1996) 499; FOGLER M. M. *et al.*, *Phys. Rev. B*, **54** (1996) 1853.
- [8] MOESSNER R. AND CHALKER J. T., *Phys. Rev. B*, **54** (1996) 5006.
- [9] PAN W. *et al.*, *Phys. Rev. Lett.*, **88** (2002) 176802.
- [10] Inter-LL excitations may be included in the random-phase approximation. This leads to a screened Coulomb interaction in a certain wave vector range but has only a minor influence on the physical properties of the system; see: ALEINER I. L. AND GLAZMAN L. I., *Phys. Rev. B*, **52** (1995) 11296.
- [11] GIRVIN S. M. *et al.*, *Phys. Rev. B*, **33** (1986) 2481.
- [12] MURTHY G. AND SHANKAR R., cond-mat/0205326 Preprint, 2002; SHANKAR R., *Phys. Rev. B*, **63** (2001) 085322.
- [13] LAM P. K. AND GIRVIN S. M., *Phys. Rev. B*, **30** (1984) 473.
- [14] ABRAHAMS E. *et al.*, *Phys. Rev. Lett.*, **42** (1979) 673.
- [15] FOGLER M. M. AND KOULAKOV A. A., *Phys. Rev. B*, **55** (1997) 9326.
- [16] SCHEIDL S. AND ROSENOW B., *Int. J. Mod. Phys. B*, **15** (2001) 1905.
- [17] MACDONALD A. H. AND FISHER M. P. A., *Phys. Rev. B*, **61** (2000) 5724.
- [18] V. OPPEN F. *et al.*, *Phys. Rev. Lett.*, **84** (2000) 2937.
- [19] GOLDMAN V. J. *et al.*, *Phys. Rev. Lett.*, **72** (1994) 2065.
- [20] LEWIS R. M. *et al.*, *Phys. Rev. Lett.*, **89** (2002) 136804.
- [21] MORF R. AND D'AMBRUMENIL N., *Phys. Rev. Lett.*, **74** (1995) 5116.
- [22] FRADKIN E. AND KIVELSON S. A., *Phys. Rev. B*, **59** (1999) 8065; WEXLER C. AND DORSEY A. T., *Phys. Rev. B*, **64** (2001) 115312.
- [23] LINDEMANN F., *Phys. Z. (Leipzig)*, **11** (1910) 69.